

<https://helda.helsinki.fi>

---

## Exploring and Reconstructing Ancestral Anatomies using Ontology-Informed Approaches

Tarasov, Sergei

2019

---

Tarasov , S , Miko , I , Yoder , M & Uyeda , J 2019 , ' Exploring and Reconstructing  
Ancestral Anatomies using Ontology-Informed Approaches ' , Biodiversity Information  
Science and Standards , vol. 3 , e37038 . <https://doi.org/10.3897/biss.3.37038>

---

<http://hdl.handle.net/10138/309613>  
<https://doi.org/10.3897/biss.3.37038>

---

cc\_by  
publishedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

Conference Abstract

# Exploring and Reconstructing Ancestral Anatomies using Ontology-Informed Approaches

Sergei Tarasov<sup>‡,§</sup>, Istvan Miko<sup>‡</sup>, Matthew Yoder<sup>¶</sup>, Josef Uyeda<sup>§</sup>

‡ Finnish Museum of Natural History, Helsinki, Finland

§ Virginia Tech, Blacksburg, United States of America

| University of New Hampshire, Durham, United States of America

¶ University of Illinois, Champaign, United States of America

Corresponding author: Sergei Tarasov ([sergxf@yandex.ru](mailto:sergxf@yandex.ru))

Received: 10 Jun 2019 | Published: 13 Jun 2019

Citation: Tarasov S, Miko I, Yoder M, Uyeda J (2019) Exploring and Reconstructing Ancestral Anatomies using Ontology-Informed Approaches. Biodiversity Information Science and Standards 3: e37038.

<https://doi.org/10.3897/biss.3.37038>

## Abstract

Ancestral character state reconstruction has been long used to gain insight into the evolution of individual traits in organisms. However, organismal anatomies (= entire phenotypes) are not merely ensembles of individual traits, rather they are complex systems where traits interact with each other due to anatomical dependencies (when one trait depends on the presence of another trait) and developmental constraints. Comparative phylogenetics has been largely lacking a method for reconstructing the evolution of entire organismal anatomies or organismal body regions. Herein, we present a new approach named *PARAMO* (*Phylogenetic Ancestral Reconstruction of Anatomy by Mapping Ontologies*, Tarasov and Uyeda 2019) that takes into account anatomical dependencies and uses stochastic maps (i.e., phylogenetic trees with an instance of mapped evolutionary history of characters, Huelsenbeck et al. 2003) along with anatomy ontologies to reconstruct organismal anatomies.

Our approach treats the entire phenotype or its component body regions as single complex characters and allows exploring and comparing phenotypic evolution at different levels of anatomical hierarchy. These complex characters are constructed by ontology-informed amalgamation of elementary characters (i.e., those coded in character matrix) using stochastic maps. In our approach, characters are linked with the terms from an anatomy

ontology, which allows viewing them not just as an ensemble of character state tokens but as entities that have their own biological meaning provided by the ontology. This ontology-informed framework provides new opportunities for tracking phenotypic radiations and anatomical evolution of organisms, which we explore using a large dataset for the insect order Hymenoptera (sawflies, wasps, ants and bees).

## Keywords

comparative phylogenetics, ontologies, organismal anatomies, trait evolution, Hymenoptera

## Presenting author

Sergei Tarasov

## Presented at

Biodiversity\_Next 2019

## References

- Huelsenbeck J, Nielsen R, Bollback J (2003) Stochastic Mapping of Morphological Characters. *Systematic Biology* 52 (2): 131-158. <https://doi.org/10.1080/10635150309342>
- Tarasov S, Uyeda J (2019) PARAMO: Phylogenetic Ancestral Reconstruction of Anatomy by Mapping Ontologies. GitHub. URL: <https://github.com/sergeitarasov/PARAMO>